arXiv:1411.5650v1 [math.AG] 20 Nov 2014

The set of fixed points of a unipotent group

Abstract. Let K be an algebraically closed field. Let G be a non-trivial connected unipotent group, which acts effectively on an affine variety X. Then every non-empty component R of the set of fixed points of G is a K-uniruled variety, i.e, there exists an affine cylinder $W \times K$ and a dominant, generically-finite polynomial mapping $\phi : W \times K \to R$. We show also that if an arbitrary infinite algebraic group G acts effectively on K^n and the set of fixed points contains a hypersurface H, then this hypersurface is K-uniruled.

1. Introduction.

Let K be an algebraically closed field (of arbitrary characteristic). Let G be a connected unipotent algebraic group, which acts effectively on a variety X. The set of fixed points of this action was studied intensively (see e.g., [**1**], [**2**], [**3**], [**4**]). In particular Białynicki-Birula has proved that if X is an affine variety, then G has no isolated fixed points.

Here we consider the case when X is an affine variety. We generalize the result of Białynicki-Birula and we prove, that the set $Fix(G)$ of fixed points of G is in fact a K-uniruled variety. In particular for every point $x \in Fix(G)$ there is a polynomial curve $\phi : K \to Fix(G)$ such that $\phi(0) = x$.

We show also that if an arbitrary infinite algebraic group G acts effectively on K^n and the set of fixed points contains a hypersurface H, then this hypersurface is K-uniruled. This generalizes [**6**].

2. Preliminaries.

At the begining we recall some basic facts about K-uniruled varieties (see [**7**]).

Proposition 2.1. *Let Γ be an affine curve. The following two statements are equivalent:*

1) there exists a regular birational map $\phi : K \to \Gamma$;

2) there exists a regular dominant map $\phi : K \to \Gamma$.

Definition 2.2. *Let Γ be an affine curve which has the property 1) (or 2)) from the above proposition. Then Γ will be called* an affine polynomial curve *and the mapping ϕ will be called* a parametrization of Γ. *A family $\mathcal{F}$ of affine polynomial curves on X is called* bounded *if there exists an embedding $i : X \subset K^N$ and a natural number D such that every curve $\Gamma \in \mathcal{F}$ has degree less then or equal to D in K^N.*

1991 *Mathematics Subject Classification.* 14L17.

Key words and phrases. affine variety, unipotent algebraic group, the set of fixed points.

The first author was partially supported by the grant of Polish Ministry of Science, 2006-2009.

REMARK 2.3. *It is easy to see that the definition of bounded family does not depend on an embedding* $i : X \to K^N$.

Now we give the definition of a K-*uniruled* variety. We have introduced this notion for uncountable fields in [**7**]. However, here we work over any field and we need a refined version of the definition (it coincides with the older one for uncountable fields).

PROPOSITION 2.4. *Let* $X \subset K^N$ *be an irreducible affine variety of dimension* ≥ 1. *The following conditions are equivalent:*

1) there is a bounded family $\mathcal{F}$ *of affine polynomial curves, such that for every point* $x \in X$ *there is a curve* $l_x \in \mathcal{F}$ *going through* x;

2) there is an open, non-empty subset $U \subset X$ *and a bounded family* $\mathcal{F}$ *of affine polynomial curves, such that for every point* $x \in U$ *there is a curve* $l_x \in \mathcal{F}$ *going through* x;

3) there exists an affine variety W *with* $\dim W = \dim X - 1$ *and a dominant polynomial mapping* $\phi : W \times K \to X$.

PROOF. 1) $\Rightarrow$ 2) is obvious. 2) $\Rightarrow$ 3) follows from [**10**]. 3) $\Rightarrow$ 2) is obvious. We prove 2) $\Rightarrow$ 1). Assume that $X \subset K^n$. Every curve $l_x \in \mathcal{F}$ is given by n polynomials of one variable:

$$l_x(t) = (x_1 + \sum_{i=1}^{D} a^{1,i}t^i, ..., x_n + \sum_{i=1}^{D} a^{n,i}t^i).$$

Let Δ denote a $nD-1$-dimensional weighted projective space with weights $1, 2, ..., D$, $..., 1, 2, ..., D$. Hence we can associate l_x with one point

$$(x_1, ..., x_n; a^{1,1}, ..., a^{1,D}; a^{2,1}, ..., a^{2,D}; ...; a^{n,1}, ..., a^{n,D}) \in X \times \Delta.$$

Let $\{f_i = 0,\ i = 1, ..., m\}$ ($f_i \in K[x_1, ..., x_n]$) be equations of the variety X. The condition $l_x \subset X$ is equivalent to conditions $f_i(l_x(t)) = 0,\ i = 1, ..., m$. The last equations are in fact equivalent to a finite number of polynomial equations

$$h_\alpha(x_1, ..., x_n; a^{1,1}, ..., a^{1,D}; a^{2,1}, ..., a^{2,D}; ...; a^{n,1}, ..., a^{n,D}) = 0,$$

which are weighted homogeneous with respect to $a^{1,1}, ..., a^{1,D}; a^{2,1}, ...; a^{n,1}, ..., a^{n,D}$. Let $W \subset X \times \Delta$ be a variety described by polynomials h_α and let $\pi : X \times \Delta \to X$ be the projection. The mapping π is proper, in particular the set $\pi(W)$ is closed. Since $\pi(W)$ contains the dense subset U, we have $\pi(W) = X$. □

Now we can state:

DEFINITION 2.5. *An affine irreducible variety* X *is called* K-uniruled *if it is of dimension* ≥ 1, *and satisfies one of equivalent conditions* 1)$-$3) *listed in Proposition 2.4.*

If the field K is uncountable we have stronger result (see [**10**]):

PROPOSITION 2.6. *Let* K *be an uncountable field. Let* X *be an irreducible affine variety of dimension* ≥ 1. *The following conditions are equivalent:*

1) X *is* K*-uniruled;*

2) for every point $x \in X$ *there is a polynomial affine curve in* X *going through* x;

3) there exists a Zariski-open, non-empty subset U *of* X, *such that for every point* $x \in U$ *there is a polynomial affine curve in* X *going through* x;

4) there exists an affine variety W with $\dim W = \dim X - 1$ *and a dominant polynomial mapping* $\phi : W \times K \to X$.

Let X be a smooth projective surface and let $D = \sum_{i=1}^{n} D_i$ be a simple normal crossing (s.n.c) divisor on X (here we consider only reduced divisors). Let $graph(D)$ be a graph of D, i.e., a graph with one vertex Q_i for each irreducible component D_i of D, and one edge between Q_i and Q_j for each point of intersection of D_i and D_j.

Definition 2.7. *Let D be a simple normal crossing divisor on a smooth surface X. We say that D is a* tree *if $graph(D)$ is connected and acyclic.*

We have the following fact which is obvious from graph theory:

Proposition 2.8. *Let X be a smooth projective surface and let divisor $D \subset X$ be a tree. Assume that $D', D'' \subset D$ are connected divisors without common components. Then D' and D'' have at most one common point.*

Definition 2.9. *Let X, Y be affine varieties and $f : X \to Y$ be a regular mapping. We say that f* is finite *at a point $y \in Y$ if there exists an open neighborhood U of y such that* $\mathrm{res}_{f^{-1}(U)} f : f^{-1}(U) \to U$ *is a finite map.*

It is well-known that if f is a generically finite, then the set of points at which f is not finite is either empty or it is a hypersurface in $\overline{f(X)}$ (for details see [**7**], [**8**]). We denote this set by S_f. Now we can formulate the following useful:

Theorem 2.10. *Let Γ be an affine curve. Let $\phi : \Gamma \times K \to K^N$ be a generically finite mapping. Then the set S_ϕ is a union of finitely many (possibly empty) of affine polynomial curves.*

Proof. Taking a normalization we can assume that the curve Γ is smooth (note that a normalization is a finite mapping). Let $\overline{\Gamma}$ be a smooth completion of Γ and denote $\overline{\Gamma} \setminus \Gamma = \{a_1, ..., a_l\}$. Let $X = \Gamma \times K$ and $\overline{X} = \overline{\Gamma} \times \mathbb{P}^1$ be a projective closure of X. The divisor $D = \overline{X} \setminus X = \overline{\Gamma} \times \infty + \sum_{i=1}^{l} \{a_i\} \times \mathbb{P}^1$ is a tree. Now we can resolve points of indeterminacy of the mapping ϕ:



Note that the divisor $D' = \pi^*(D)$ is a tree. Let $\overline{\Gamma} \times \infty'$ denote a proper transform of $\overline{\Gamma} \times \infty$. It is an easy observation that $\phi'(\overline{\Gamma} \times \infty') \subset H_\infty$, where H_∞ denotes the hyperplane at infinity of $\mathbb{P}^N$. Now $S_\phi = \phi'(D' \setminus \phi'^{-1}(H_\infty))$. The curve $L = \phi'^{-1}(H_\infty)$

is a complement of a semi-affine variety $\phi'^{-1}(K^N)$ hence it is connected (for details see [**7**], Lemma 4.5). Now by Proposition 2.8 we have that every irreducible curve $Z \subset D'$ which does not belong to L has at most one common point with L. Let $S \subset S_\phi$ be an irreducible component. Hence S is a curve. There is a curve $Z \subset D'$, which has exactly one common point with L such that $S = \phi'(Z \setminus L) = \phi'(K)$. This completes the proof. □

3. Main Result.

The aim of this section is to prove the following:

THEOREM 3.1. *Let G be a non-trivial connected unipotent group which acts effectively on an affine variety X. Then every non-empty component R of the set of fixed points of G is a K-uniruled variety.*

PROOF. First of all let us recall that a connected unipotent group has a normal series

$$0 = G_0 \subset G_1 \subset ... \subset G_r = G,$$

where $G_i/G_{i-1} \cong G_a = (K, +, 0)$. By induction on $\dim G$ we can easily reduce the general case to that of $G = G_a$.

First assume that the field K is uncountable. Take a point $a \in R$. By Proposition 2.6 it is enough to prove, that there is an affine polynomial curve $S \subset Fix(G)$ through a. Let L be an irreducible curve in X going through a, which is not contained in any orbit of G and it is not contained in $Fix(G)$. Consider a surface $Y = L \times G$. There is natural G action on Y : for $h \in G$ and $y = (l, g) \in Y$ we put $h(y) = (l, hg) \in Y$. Take a mapping

$$\Phi : L \times G \ni (x, g) \to g(x) \in X.$$

It is a generically-finite polynomial mapping. Observe that it is G-invariant i.e., $\Phi(gy) = g\Phi(y)$. This implies that the set S_Φ of points at which the mapping Φ is not finite is G-invariant. Indeed, it is enough to show that the complement of this set is G-invariant. Let Φ be finite at $x \in X$. This means that there is an open neighborhood U of x such that the mapping $\Phi : \Phi^{-1}(U) \to U$ is finite. Now we have the following diagram:

$$\begin{array}{ccc} \Phi^{-1}(U) & \xrightarrow{\ g\ } & \Phi^{-1}(gU) = g\Phi^{-1}(U) \\ \downarrow \Phi & & \downarrow \Phi \\ U & \xrightarrow{\ g\ } & gU \end{array}$$

This diagram shows that the mapping Φ is finite over gU if it is finite over U. In particular this implies that the set S_Φ is G-invariant. Let $S_\Phi = S_1 \cup S_2 ... \cup S_k$ be a decomposition of S_Φ in (irreducible) affine polynomial curves (see Theorem 2.10). Since the set S_Φ is G-invariant, we have that each curve S_i is also G-invariant.

Note that the point a belongs to S_Φ, because the fiber over a has infinite number of points. We can assume that $a \in S_1$. Let $x \in S_1$, we want to show that $x \in Fix(G)$. Indeed, otherwise $G.x = S_1$ and a would be in the orbit of x - a contradiction. Hence $S_1 \subset Fix(G)$ and we conclude our result by Theorem 2.10.

Now assume that the field K is countable. Let $X \subset K^n$. Let T be uncountable algebraically closed extension of K. By the base change the group G acts on $\overline{X} \subset T^n$. Moreover, the variety $\overline{R} \subset T^n$ is a component of the set of fixed points of G (because the set R is dense in $\overline{R}$). By the first part of our proof the variety $\overline{R}$ is T-uniruled. In particular there exists a number D such that for every point $x \in \overline{R}$ there is a polynomial affine curve $l_x \subset \overline{R} \subset T^n$, of degree at most D, going through x. Note that it is true for every point $x \in R$.

Every such curve l_x is given by n polynomials of one variable:

$$l_x(t) = (x_1 + \sum_{i=1}^{d} a^{1,i}t^i, ..., x_n + \sum_{i=1}^{d} a^{n,i}t^i),$$

where $d \leq D$. Hence we can associate l with one point

$$(a^{1,0}, a^{1,1}, ..., a^{1,d}; a^{2,0}, ..., a^{2,d}; ...; a^{n,0}, ..., a^{n,d}) \in T^n.$$

We can assume without loss of generality that $a^1_d = 1$. Let $\{f_i = 0,\ i = 1, ..., m\}$ ($f_i \in K[x_1, ..., x_n]$) be equations of the variety S. The condition $l_x \subset \overline{R}$ is equivalent to conditions $f_i(l(t)) = 0,\ i = 1, ..., m$. The last equations are in fact equivalent to a finite number of polynomial equations

$$h_\alpha(a^{1,0}, a^{1,1}, ..., a^{1,d}; a^{2,0}, ..., a^{2,d}; ...; a^{n,0}, ..., a^{n,d}) = 0,$$

where $h_\alpha \in K[y_1, ..., y_N]$. Equations $h_\alpha = 0$ plus extra conditions $a^{i,0} = x_i,\ i = 1, ..., n$ and $a^{1,d} = 1$ have solutions in the field T, hence they have also solutions in the field K.

This means that we can find an affine polynomial curve l_x over the field K of degree at most D, which is contained in R and goes through x. Consequently the variety R is K-uniruled. The proof of Theorem 3.1 is complete. □

COROLLARY 3.2. *(Białynicki-Birula, [**1**]). Let G be a non-trivial connected unipotent group which acts effectively on an affine variety X. Then G has no isolated fixed points.*

Theorem 3.1 (or rather its proof) suggests a following generalization of [**6**]:

THEOREM 3.3. *Let G be an infinite connected algebraic group which acts effectively on K^n, $n \geq 2$. Assume that an irreducible hypersurface W is contained in the set of fixed points of G. Then W is K- uniruled.*

PROOF. Since G acts effectively on affine space K^n we can assume by the Chevalley Theorem (see [**9**], Th. C, p.190) that the group G is affine. In particular it contains either the subgroup $G_m = (K^*, \cdot, 1)$ or the subgroup $G_a = (K, +, 0)$ (see e.g., [**5**]). Thus we can assume that G is either G_m or it is G_a.

As before we can assume that the field K is uncountable. Take a point $a \in W$. By Proposition 2.6 it is enough to prove, that there is an affine parametric curve $S \subset W$ through a. Let L be a line in K^n going through a such that the set $L \cap Fix(G)$ is finite. Set $L \cap W = \{a, a_1, ..., a_m\}$. Now consider a mapping

$$\phi : L \times G \ni (x, g) \to g(x) \in K^n.$$

Observe that $\phi(L\times G)$ is a union of disjoint orbits of G. This implies $\phi(L\times G)\cap W = \{a, a_1, ..., a_m\}$. Take $X = \overline{\phi(L\times G)}$. Note that $X\cap W$ is a union of curves. This means that there is a curve $S \subset X\cap W$, which contains the point a. However $S \subset \overline{X\setminus\phi(L\times G)}$. This implies that $S \subset S_\phi$ and we conclude by Theorem 2.10. $\square$

To finish this note we state:

Conjecture. *Let K be an algebraically closed field. Let G be an algebraic group, which acts effectively on K^n. If S is an irreducible component of the set of fixed points of G, then S is either a point or it is a K-uniruled variety.*

Acknowledgments

We are grateful to Andrzej Białynicki-Birula for helpful discussions. We also want to thank Tadeusz Mostowski for helpful remarks.

References

[1] A. Białynicki-Birula, On fixed point schemes of actions of multiplicative and additive groups, Topology 12 (1973), 99-103.
[2] J. Carrell, A. Sommese, A generalization theorem of Horrocks, in: Group Action and Vector Fields, in: Lecture Notes in Math. 956, Springer-Verlag 1982.
[3] D. Gross, On the fundamental group of the fixed points of an unipotent actions, Manuscripta Math. 60 (1988), 487-496.
[4] G. Horrocks, Fixed point schemes of actions of additive group actions, Topology 8 (1969), 233-242.
[5] J. Humphreys, Linear Algebraic Groups, Springer-Verlag 1998.
[6] Z. Jelonek, The set of fixed points of an algebraic group, Bull. Acad. Polon. Sci. Math. 51 (2003), 69-73.
[7] Z. Jelonek, Testing sets for properness of polynomial mappings, Math. Ann. 315 (1999), 1-35.
[8] Z. Jelonek, Topological characterization of finite mappings, Bull. Acad. Polon. Sci. Math. 49 (2001), 279-283.
[9] I.R. Shafarevich, Basic Algebraic Geometry, Springer-Verlang, 1974.
[10] A. Stasica, Geometry of the Jelonek set, J. Pure Appl. Algebra 137 (1999), 49-55.

(Z. Jelonek) Instytut Matematyki, Polska Akademia Nauk, Św. Tomasza 30, 31-027 Kraków, Poland
E-mail address: najelone@cyf-kr.edu.pl

(M. Lasoń) Instytut Matematyki, Polska Akademia Nauk, Św. Tomasza 30, 31-027 Kraków, Poland; Wydział Matematyki i Informatyki, Uniwersytet Jagielloński, Prof. S. Łojasiewicza 6, 30-348 Kraków, Poland
E-mail address: michalason@gmail.com